

Revision Date: 11/23/2020





PRODUCT SPECIFICATION

Skim Milk Powder (SMP) LH Instantized

Valley Milk, LLC Skim Milk Powder Low Heat Instantized is made through the spray ball method of drying fresh skimmed, pasteurized cow's milk and incorporating a soy based lecithin during the process. All milk that is processed in our facility is collected from our dedicated multigenerational, family owned dairy farms. All of the milk originates from the United States in California's San Joaquin Valley and Grade A SMP Low Heat Instantized has a shelf life of two years from the production date.

PHYSICAL

	MINIMUM	MAXIMUM	UNITS	METHODS
				AOAC
Protein	32.40		%m/m	991.20
Milk Protein in				AOAC
milk Solids Non-				991.20
Fat	34.0	-	%m/m	
Casein Protein	27.0	-	%m/m	AOAC
Moisture	-	4.0	%m/m	AOAC 927.05
Fat	-	1.25	%m/m	AOAC 989.05
				ADPI/Std. M. 17th
Titratable acidity	-	0.15	%m/v	ed.
				ADPI/Std. M. 17th
WPNI	6.0	-	mg/g	ed.
Lactose	-	58.3	%m/v	AOAC 984.22
Ash	-	8.6	%m/v	AOAC 945.46
				ADPI/Std. M. 17th
рН	6.4	7.2	mol/L	ed.
				ADPI/Std. M. 17th
Solubility Index	-	1.25	mL	ed.
		Not		
Foreign matter	-	Detected	/25g	
Scorched			mg/25	ADPI BUL. 916
particles	-	7.5	g	

CHEMICAL

	MINIMUM	MAXIMUM	UNITS	METHODS
Antibiotics	-	Negative	Charm S/L	Charm SL3
Aflatoxin M1	-	<0.50	μg/L	Charm

SENSORY

Flavor	Free from objectionable flavors	
Appearance	No notable defect in color or particle	
Color	White to light cream	

MICROBIOLOGICAL

	MINIMUM	MAXIMUM	UNITS	METHODS
Aerobic Plate Count	-	<10,000	cfu/g	AOAC 990.12
Yeast & Mold	-	<50	cfu/g	BAM Ch 18
Coliforms	-	<10	cfu/g	AOAC 991.14
Escherichia coli	-	<3	MPN/g	BAM Ch 4
Coagulase- Positive Staphylococci	-	Negative	cfu/g	AOAC 2003.08
Salmonella	-	Absent	/375g	AOAC-RI 121501
Listeria	-	Negative	/25g	AOAC-RI 061702



PACKAGING

Product may be packaged in either heat sealed poly-lined multiwall kraft paper bags or packaged into poly lined totes without the use of staples or metal fasters.



STORAGE AND HANDLING

It is recommended that product is stored at 80°F or less, with relative humidity of 65% or less, and in an odor-free environment. Stocks should be used in rotation preferably within 24 months of manufacture or as dictated by the expiration date labeled on the product.